



Multiple Polylogarithms: An Introduction

Michel Waldschmidt

► To cite this version:

Michel Waldschmidt. Multiple Polylogarithms: An Introduction. Conference on number theory and discrete mathematics in honour of Srinivasa Ramanujan, Oct 2000, Chandigarh, India. hal-00416166

HAL Id: hal-00416166

<https://hal.science/hal-00416166>

Submitted on 12 Sep 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Multiple Polylogarithms: An Introduction

by

M. WALDSCHMIDT(*)

Abstract

Multiple polylogarithms in a single variable are defined by

$$\mathrm{Li}_{(s_1, \dots, s_k)}(z) = \sum_{n_1 > n_2 > \dots > n_k \geq 1} \frac{z^{n_1}}{n_1^{s_1} \dots n_k^{s_k}},$$

when s_1, \dots, s_k are positive integers and z a complex number in the unit disk. For $k = 1$, this is the classical polylogarithm $\mathrm{Li}_s(z)$. These multiple polylogarithms can be defined also in terms of iterated Chen integrals and satisfy *shuffle relations*. Multiple polylogarithms in several variables are defined for $s_i \geq 1$ and $|z_i| < 1$ ($1 \leq i \leq k$) by

$$\mathrm{Li}_{(s_1, \dots, s_k)}(z_1, \dots, z_k) = \sum_{n_1 > n_2 > \dots > n_k \geq 1} \frac{z_1^{n_1} \dots z_k^{n_k}}{n_1^{s_1} \dots n_k^{s_k}},$$

and they satisfy not only shuffle relations, but also *stuffle relations*. When one specializes the shuffle relations in one variable at $z = 1$ and the stuffle relations in several variables at $z_1 = \dots = z_k = 1$, one gets linear or quadratic dependence relations between the Multiple Zeta Values

$$\zeta(s_1, \dots, s_k) = \sum_{n_1 > n_2 > \dots > n_k \geq 1} \frac{1}{n_1^{s_1} \dots n_k^{s_k}}$$

which are defined for k, s_1, \dots, s_k positive integers with $s_1 \geq 2$. The *Main Diophantine Conjecture* states that one obtains in this way all algebraic relations between these MZV.

Classification AMS: 11J91, 33E30

(*) Institut de Mathématiques, Université P. et M. Curie (Paris VI),
Théorie des Nombres Case 247, F-75013 PARIS miw@math.jussieu.fr
<http://www.math.jussieu.fr/~miw/>

0. Introduction

A long term project is to determine all algebraic relations among the values

$$\pi, \zeta(3), \zeta(5), \dots, \zeta(2n+1), \dots$$

of the Riemann zeta function

$$\zeta(s) = \sum_{n \geq 1} \frac{1}{n^s}.$$

So far, one only knows that the first number in this list, π , is transcendental, that the second one, $\zeta(3)$, is irrational, and that the other ones span a \mathbb{Q} -vector space of infinite dimension [R], [BR].

The expected answer is disappointingly simple: it is widely believed that there are no relations, which means that these numbers should be algebraically independent:

(?) *For any $n \geq 0$ and any nonzero polynomial $P \in \mathbb{Z}[X_0, \dots, X_n]$,*

$$P(\pi, \zeta(3), \zeta(5), \dots, \zeta(2n+1)) \neq 0.$$

If true, this property would mean that there is no interesting algebraic structure.

The situation changes drastically if we enlarge our set so as to include the so-called Multiple Zeta Values (MZV, also called Euler-Zagier numbers – see [Eu] and [Z]):

$$\zeta(s_1, \dots, s_k) = \sum_{n_1 > n_2 > \dots > n_k \geq 1} \frac{1}{n_1^{s_1} \dots n_k^{s_k}},$$

which are defined for k, s_1, \dots, s_k positive integers with $s_1 \geq 2$. It may be hoped that the initial goal could be reached if one could determine all algebraic relations between the MZV. Now there are plenty of relations between them, providing a rich algebraic structure. One type of such relations arises when one multiplies two such series: it is easy to see that one gets a linear combination of MZV. There is another type of algebraic relations between MZV, coming from their expressions as integrals. Again the product of two such integrals is a linear combination of MZV. Following [B³], we will use the name *stuffle* for the relations arising from the series, and *shuffle* for those arising from the integrals.

The *Main Diophantine Conjecture* (Conjecture 5.3 below) states that these relations are sufficient to describe all algebraic relations between MZV. One should be careful when stating such a conjecture: it is necessary to include some relations which are deduced from the stuffle and shuffle applied to divergent series (i.e. with $s_1 = 1$).

There are several ways of dealing with the divergent case. Here, we use the *multiple polylogarithms*

$$\text{Li}_{(s_1, \dots, s_k)}(z) = \sum_{n_1 > n_2 > \dots > n_k \geq 1} \frac{z^{n_1}}{n_1^{s_1} \dots n_k^{s_k}},$$

which are defined for $|z| < 1$ when s_1, \dots, s_k are all ≥ 1 , and which are also defined for $|z| = 1$ if $s_1 \geq 2$.

These multiple polylogarithms can be expressed as iterated Chen integrals, and from this representation one deduces shuffle relations. There is no stuffle relations for multiple polylogarithms in a single variable, but one recovers them by introducing the multivariable functions

$$\text{Li}_{(s_1, \dots, s_k)}(z_1, \dots, z_k) = \sum_{n_1 > n_2 > \dots > n_k \geq 1} \frac{z_1^{n_1} \dots z_k^{n_k}}{n_1^{s_1} \dots n_k^{s_k}} \quad (s_i \geq 1, 1 \leq i \leq k)$$

which are defined not only for $|z_i| < 1$ ($1 \leq i \leq k$), but also for $|z_i| \leq 1$ ($1 \leq i \leq k$) if $s_1 \geq 2$.

Notation. Given a string a_1, \dots, a_k of integers, the notation $\{a_1, \dots, a_k\}_n$ stands for the kn -tuple

$$(a_1, \dots, a_k, \dots, a_1, \dots, a_k),$$

where the string a_1, \dots, a_k is repeated n times.

1. Multiple Polylogarithms in One Variable and Multiple Zeta Values

Let k, s_1, \dots, s_k be positive integers. Write \underline{s} in place of (s_1, \dots, s_k) . One defines a complex function of one variable by

$$\text{Li}_{\underline{s}}(z) = \sum_{n_1 > n_2 > \dots > n_k \geq 1} \frac{z^{n_1}}{n_1^{s_1} \dots n_k^{s_k}}.$$

This function is analytic in the open unit disk, and, in the case $s_1 \geq 2$, it is also continuous on the closed unit disk. In the latter case we have

$$\zeta(\underline{s}) = \text{Li}_{\underline{s}}(1).$$

One can also define in an equivalent way these functions by induction on the number $p = s_1 + \dots + s_k$ (the *weight* of \underline{s}) as follows. Plainly we have

$$(1.1) \quad z \frac{d}{dz} \text{Li}_{(s_1, \dots, s_k)}(z) = \text{Li}_{(s_1-1, s_2, \dots, s_k)}(z) \quad \text{if} \quad s_1 \geq 2$$

and

$$(1.2) \quad (1-z) \frac{d}{dz} \text{Li}_{(1,s_2,\dots,s_k)}(z) = \text{Li}_{(s_2,\dots,s_k)}(z).$$

Together with the initial conditions

$$(1.3) \quad \text{Li}_{\underline{s}}(0) = 0,$$

the differential equations (1.1) and (1.2) determine all the $\text{Li}_{\underline{s}}$.

Therefore, as observed by M. Kontsevich (cf. [Z]; see also [K] Chap. XIX, § 11 for an early reference to H. Poincaré, 1884), an equivalent definition for $\text{Li}_{\underline{s}}$ is given by integral formulae as follows. Starting^(*) with $k = s = 1$, we write

$$\text{Li}_1(z) = -\log(1-z) = \int_0^z \frac{dt}{1-t},$$

where the complex integral is over any path from 0 to z inside the unit circle. From the differential equations (1.1) one deduces, by induction, for $s \geq 2$,

$$\text{Li}_s(z) = \int_0^z \text{Li}_{s-1}(t) \frac{dt}{t} = \int_0^z \frac{dt_1}{t_1} \int_0^{t_1} \frac{dt_2}{t_2} \cdots \int_0^{t_{s-2}} \frac{dt_{s-1}}{t_{s-1}} \int_0^{t_{s-1}} \frac{dt_s}{1-t_s}.$$

In the last formula, the complex integral which is written on the left (and which is the last to be computed) is over any path inside the unit circle from 0 to z , the second one is from 0 to t_1, \dots and the last one on the right, which is the first to be computed, is from 0 to t_{s-1} .

Then iterated integrals (see [K] Chap. XIX, § 11) provide a compact form for such expressions as follows. For $\varphi_1, \dots, \varphi_p$ differential forms and x, y complex numbers, define inductively

$$\int_x^y \varphi_1 \cdots \varphi_p = \int_x^y \varphi_1(t) \int_x^t \varphi_2 \cdots \varphi_p.$$

For $\underline{s} = (s_1, \dots, s_k)$, set

$$\omega_{\underline{s}} = \omega_0^{s_1-1} \omega_1 \cdots \omega_0^{s_k-1} \omega_1,$$

where

$$\omega_0(t) = \frac{dt}{t} \quad \text{and} \quad \omega_1(t) = \frac{dt}{1-t}.$$

(*) This induction could as well be started from $k = 0$, provided that we set $\text{Li}_{\emptyset}(z) = 1$.

Then the differential equations (1.1) and (1.2) with initial conditions (1.3) can be written

$$(1.4) \quad \text{Li}_{\underline{s}}(z) = \int_0^z \omega_{\underline{s}}.$$

Example. For any $n \geq 1$ and $|z| < 1$ we have

$$(1.5) \quad \text{Li}_{\{1\}_n}(z) = \frac{1}{n!} \left(\log(1/(1-z)) \right)^n,$$

which can be written in terms of generating series as

$$\sum_{n=0}^{\infty} \text{Li}_{\{1\}_n}(z) x^n = (1-z)^{-x}.$$

The constant term $\text{Li}_{\{1\}_0}(z)$ is 1.

2. Shuffle Product and the First Standard Relations

Denote by $X = \{\omega_0, \omega_1\}$ the alphabet with two letters and by X^* the set of words on X . A word is nothing else than a non-commutative monomial in the two letters ω_0 and ω_1 . The linear combinations of such words with rational coefficients

$$\sum_u c_u u,$$

where $\{c_u ; u \in X^*\}$ is a set of rational numbers with finite support, is the non-commutative ring $\mathbb{Q}\langle\omega_0, \omega_1\rangle$. We are interested with the set $X^*\omega_1$ of words which end with ω_1 , together with the empty word \emptyset . The linear combinations of such words is a left ideal of $\mathbb{Q}\langle\omega_0, \omega_1\rangle$ which we denote by $\mathbb{Q}\langle\omega_0, \omega_1\rangle\omega_1$.

The set $X^*\omega_1$ is also the set of words $\omega_{\underline{s}}$, with $\underline{s} = (s_1, \dots, s_k)$. We define $\text{Li}_u(z)$ for $u \in X^*\omega_1$ by $\text{Li}_u(z) = \text{Li}_{\underline{s}}(z)$ when $u = \omega_{\underline{s}}$. By linearity we extend the definition of $\text{Li}_u(z)$ to the ideal $\mathbb{Q}\langle\omega_0, \omega_1\rangle\omega_1$:

$$\text{Li}_v(z) = \sum_u c_u \text{Li}_u(z) \quad \text{for} \quad v = \sum_u c_u u.$$

The set of *convergent words* is the set, denoted by $\omega_0 X^* \omega_1$, of words which start with ω_0 and end with ω_1 (including the empty word). The \mathbb{Q} -vector subspace they span in $\mathbb{Q}\langle\omega_0, \omega_1\rangle$ is denoted by $\omega_0 \mathbb{Q}\langle\omega_0, \omega_1\rangle \omega_1$, and for v in this space we set

$$\zeta(v) = \text{Li}_v(1).$$

Definition. The shuffle product of two words in X^* is the element in $\mathbb{Q}\langle\omega_0, \omega_1\rangle$ which is defined inductively as follows:

$$\emptyset \sqcup u = u \sqcup \emptyset = u$$

for any u in $X^*\omega_1$, and

$$(\omega_i u) \sqcup (\omega_j v) = \omega_i (u \sqcup \omega_j v) + \omega_j (\omega_i u \sqcup v)$$

for u, v in X^* and i, j equal to 0 or 1.

This product is extended bilinearly to $\mathbb{Q}\langle\omega_0, \omega_1\rangle$ and defines a commutative and associative law. Moreover $\mathbb{Q}\langle\omega_0, \omega_1\rangle\omega_1$ is stable under \sqcup .

Computing the product $\text{Li}_u(z)\text{Li}_{u'}(z)$ of the two associated Chen iterated integrals yields (see [MPH], Th. 2):

Proposition 2.1. For u and u' in $X^*\omega_1$,

$$\text{Li}_u(z)\text{Li}_{u'}(z) = \text{Li}_{u \sqcup u'}(z).$$

For instance from

$$\omega_1 \sqcup \omega_0 \omega_1 = \omega_1 \omega_0 \omega_1 + 2\omega_0 \omega_1^2$$

we deduce

$$(2.2) \quad \text{Li}_1(z)\text{Li}_2(z) = \text{Li}_{1,2}(z) + 2\text{Li}_{2,1}(z).$$

Setting $z = 1$, we deduce from Proposition 2.1:

$$(2.3) \quad \zeta(u)\zeta(u') = \zeta(u \sqcup u')$$

for u and u' in $\omega_0 X^*\omega_1$.

These are the *first standard relations* between multiple zeta values.

3. Shuffle Product for Multiple Polylogarithms in Several Variables

The functions of k complex variables^(*)

$$\text{Li}_{\underline{s}}(z_1, \dots, z_k) = \sum_{n_1 > n_2 > \dots > n_k \geq 1} \frac{z_1^{n_1} \dots z_k^{n_k}}{n_1^{s_1} \dots n_k^{s_k}}$$

(*) Our notation for

$$\text{Li}_{(s_1, \dots, s_k)}(z_1, \dots, z_k)$$

is the same as in [W], but it corresponds to Goncharov's notation [G2] for

$$\text{Li}_{(s_k, \dots, s_1)}(z_k, \dots, z_1).$$

have been considered as early as 1904 by N. Nielsen, and rediscovered later by A.B. Goncharov [G1,G2]. Recently, J. Écalle [É] used them for z_i roots of unity (in case $s_1 \geq 2$): these are the *decorated multiple polylogarithms*. Of course one recovers the one variable functions $\text{Li}_{\underline{s}}(z)$ by specializing $z_2 = \dots = z_k = 1$. For simplicity we write $\text{Li}_{\underline{s}}(\underline{z})$, where \underline{z} stands for (z_1, \dots, z_k) . There is an integral formula which extends (1.4). Define

$$\omega_z(t) = \begin{cases} \frac{zdt}{1-zt} & \text{if } z \neq 0, \\ \frac{dt}{t} & \text{if } z = 0. \end{cases}$$

From the differential equations

$$z_1 \frac{\partial}{\partial z_1} \text{Li}_{\underline{s}}(\underline{z}) = \text{Li}_{(s_1-1, s_2, \dots, s_k)}(\underline{z}) \quad \text{if } s_1 \geq 2$$

and

$$(1 - z_1) \frac{\partial}{\partial z_1} \text{Li}_{(1, s_2, \dots, s_k)}(\underline{z}) = \text{Li}_{(s_2, \dots, s_k)}(z_1 z_2, z_3, \dots, z_k),$$

generalizing (1.1) and (1.2), we deduce

$$\text{Li}_{\underline{s}}(\underline{z}) = \int_0^1 \omega_0^{s_1-1} \omega_{z_1} \omega_0^{s_2-1} \omega_{z_1 z_2} \dots \omega_0^{s_k-1} \omega_{z_1 \dots z_k}.$$

Because of the occurrence of the products $z_1 \dots z_j$ ($1 \leq j \leq k$), the authors of [G1] and [B³L] perform the change of variables

$$y_j = z_1^{-1} \dots z_j^{-1} \quad (1 \leq j \leq k) \quad \text{and} \quad z_j = \frac{y_{j-1}}{y_j} \quad (1 \leq j \leq k)$$

with $y_0 = 1$, and introduce the differential forms

$$\omega'_y(t) = -\omega_{y^{-1}}(t) = \frac{dt}{t-y},$$

so that $\omega'_0 = \omega_0$ and $\omega'_1 = -\omega_1$. Also they define

$$\begin{aligned} \lambda \left(\begin{matrix} s_1, \dots, s_k \\ y_1, \dots, y_k \end{matrix} \right) &= \text{Li}_{\underline{s}}(1/y_1, y_1/y_2, \dots, y_{k-1}/y_k) \\ &= \sum_{\nu_1 \geq 1} \dots \sum_{\nu_k \geq 1} \prod_{j=1}^k y_j^{-\nu_j} \left(\sum_{i=j}^k \nu_i \right)^{-s_j} \\ &= (-1)^p \int_{\Delta_p} \omega_0^{s_1-1} \omega'_{y_1} \dots \omega_0^{s_k-1} \omega'_{y_k}. \end{aligned}$$

With this notation some formulae are simpler. For instance the shuffle relation is easier to write with λ : the shuffle is defined on words on the alphabet $\{\omega'_y ; y \in \mathbb{C}\}$, (including $y = 0$), inductively by

$$(\omega'_y u) \sqcup (\omega'_{y'} v) = \omega'_y (u \sqcup \omega'_{y'} v) + \omega'_{y'} (\omega'_y u \sqcup v).$$

4. Stuffle Product and the Second Standard Relations

The functions $\text{Li}_{\underline{s}}(\underline{z})$ satisfy not only shuffle relations, but also *stuffle relations* arising from the product of two series:

$$(4.1) \quad \text{Li}_{\underline{s}}(\underline{z})\text{Li}_{\underline{s}'}(\underline{z}') = \sum_{\underline{s}''} \text{Li}_{\underline{s}''}(\underline{z}''),$$

where the notation is as follows: \underline{s}'' runs over the tuples $(s''_1, \dots, s''_{k''})$ obtained from $\underline{s} = (s_1, \dots, s_k)$ and $\underline{s}' = (s'_1, \dots, s'_{k'})$ by inserting, in all possible ways, some 0 in the string (s_1, \dots, s_k) as well as in the string $(s'_1, \dots, s'_{k'})$ (including in front and at the end), so that the new strings have the same length k'' , with $\max\{k, k'\} \leq k'' \leq k + k'$, and by adding the two sequences term by term. For each such \underline{s}'' , the component z''_i of \underline{z}'' is z_j if the corresponding s''_i is just s_j (corresponding to a 0 in \underline{s}'), it is z'_ℓ if the corresponding s''_i is s'_ℓ (corresponding to a 0 in \underline{s}), and finally it is $z_j z'_\ell$ if the corresponding s''_i is $s_j + s'_\ell$. For instance

\underline{s}	s_1	s_2	0	s_3	s_4	\dots	0
\underline{s}'	0	s'_1	s'_2	0	s'_3	\dots	$s'_{k'}$
\underline{s}''	s_1	$s_2 + s'_1$	s'_2	s_3	$s_4 + s'_3$	\dots	$s'_{k'}$
\underline{z}''	z_1	$z_2 z'_1$	z'_2	z_3	$z_4 z'_3$	\dots	$z'_{k'}$

Of course the 0's are inserted so that no s''_i is zero.

Examples. For $k = k' = 1$ the stuffle relation (4.1) yields

$$(4.2) \quad \text{Li}_s(z)\text{Li}_{s'}(z') = \text{Li}_{(s,s')}(z, z') + \text{Li}_{(s',s)}(z', z) + \text{Li}_{s+s'}(zz'),$$

while for $k = 1$ and $k' = 2$ we have

$$(4.3) \quad \begin{aligned} \text{Li}_s(z)\text{Li}_{(s'_1, s'_2)}(z'_1, z'_2) &= \text{Li}_{(s, s'_1, s'_2)}(z, z'_1, z'_2) + \text{Li}_{(s'_1, s, s'_2)}(z'_1, z, z'_2) \\ &+ \text{Li}_{(s'_1, s'_2, s)}(z'_1, z'_2, z) + \text{Li}_{(s+s'_1, s'_2)}(zz'_1, z'_2) + \text{Li}_{(s'_1, s+s'_2)}(z'_1, zz'_2). \end{aligned}$$

The *stuffle product* is defined on $X^*\omega_1$ inductively by

$$\emptyset * u = u * \emptyset = u$$

for $u \in X^*\omega_1$ and

$$\begin{aligned} (\omega_0^{s-1}\omega_1 u) * (\omega_0^{t-1}\omega_1 u') &= \\ \omega_0^{s-1}\omega_1(u * \omega_0^{t-1}\omega_1 u') &+ \omega_0^{t-1}\omega_1(\omega_0^{s-1}\omega_1 u * u') + \omega_0^{s+t-1}\omega_1(u * u') \end{aligned}$$

for u and u' in $X^*\omega_1$, $s \geq 1$, $t \geq 1$.

Specializing (4.1) at $z_1 = \cdots = z_k = z'_1 = \cdots = z'_{k'} = 1$, we deduce

$$(4.4) \quad \zeta(u)\zeta(u') = \zeta(u * u')$$

for u and u' in $\omega_0 X^* \omega_1$.

These are the *second standard relations* between multiple zeta values. For instance (4.3) with $z = z'_1 = z'_2 = 1$ gives

$$\begin{aligned} \zeta(s)\zeta(s'_1, s'_2) &= \zeta(s, s'_1, s'_2) + \zeta(s'_1, s, s'_2) + \zeta(s'_1, s'_2, s) \\ &\quad + \zeta(s + s'_1, s'_2) + \zeta(s'_1, s + s'_2) \end{aligned}$$

for $s \geq 2$, $s'_1 \geq 2$ and $s'_2 \geq 1$.

5. The Third Standard Relations and the Main Diophantine Conjectures

We start with an example. Combining the stuffle relation (4.2) for $s = s' = 1$ with the shuffle relation (2.2) for $z' = z$, we deduce

$$(5.1) \quad \text{Li}_{1,2}(z, 1) + 2\text{Li}_{2,1}(z, 1) = \text{Li}_{1,2}(z, z) + \text{Li}_{2,1}(z, z) + \text{Li}_3(z^2).$$

The two sides are analytic inside the unit circle, but not convergent at $z = 1$. We claim that

$$F(z) = \text{Li}_{1,2}(z, 1) - \text{Li}_{1,2}(z, z) = \sum_{n_1 > n_2 \geq 1} \frac{z^{n_1}(1 - z^{n_2})}{n_1 n_2^2}$$

tends to 0 as z tends to 1 inside the unit circle. Indeed for $|z| < 1$ we have

$$|1 - z^{n_2}| = |(1 - z)(1 + z + \cdots + z^{n_2-1})| < n_2 |1 - z|,$$

hence

$$\sum_{n_2=1}^{n_1-1} \frac{|1 - z^{n_2}|}{n_2^2} < |1 - z| \sum_{n_2=1}^{n_1-1} \frac{1}{n_2}.$$

From (1.5) with $n = 2$ we deduce

$$|F(z)| \leq |1 - z| \text{Li}_{1,1}(|z|) = \frac{1}{2} |1 - z| \left(\log(1/(1 - |z|)) \right)^2.$$

Therefore, taking the limit of the relation (5.1) as $z \rightarrow 1$ yields Euler's formula

$$\zeta(2, 1) = \zeta(3).$$

This argument works in a quite general setting and yields the relations

$$(5.2) \quad \zeta(\omega_1 * u - \omega_1 \sqcup u) = 0$$

for each $u \in \omega_0 X^* \omega_1$.

These are the *third standard relations* between multiple zeta values.

Zagier, Goncharov, Kontsevich, ... (see [Z]) conjecture that the three standard relations (2.3), (4.4) and (5.2) constitute a basis of the ideal of algebraic relations between all numbers $\zeta(\underline{s})$. Here are precise statements.

We introduce independent variables Z_u , where u ranges over the set $X^* \omega_1$. For $v = \sum_u c_u u$ in $\mathbb{Q}\langle \omega_0, \omega_1 \rangle \omega_1$, we set

$$Z_v = \sum_u c_u Z_u.$$

In particular for u_1 and u_2 in $\omega_0 X^* \omega_1$, $Z_{u_1 \sqcup u_2}$ and $Z_{u_1 * u_2}$ are linear forms in Z_u , $u \in \omega_0 X^* \omega_1$. Also, for $v \in \omega_0 \mathbb{Q}\langle \omega_0, \omega_1 \rangle \omega_1$, $Z_{\omega_1 \sqcup v - \omega_1 * v}$ is a linear form in Z_u , $u \in \omega_0 X^* \omega_1$.

Denote by R the ring of polynomials with coefficients in \mathbb{Q} in the variables Z_u where u ranges over the set $\omega_0 X^* \omega_1$, and by \mathfrak{I} the ideal of R consisting of all polynomials which vanish under the specialization map

$$Z_u \mapsto \zeta(u) \quad (u \in \omega_0 X^* \omega_1).$$

Conjecture 5.3. *The polynomials*

$$Z_u Z_v - Z_{u \sqcup v}, \quad Z_u Z_v - Z_{u * v} \quad \text{and} \quad Z_{\omega_1 \sqcup u - \omega_1 * u},$$

where u and v range over the set of elements in $\omega_0 X^* \omega_1$, generate the ideal \mathfrak{I} .

Denote by \mathfrak{Z}_p the \mathbb{Q} -vector subspace of \mathbb{R} spanned by the real numbers $\zeta(\underline{s})$ with \underline{s} of weight p , with $\mathfrak{Z}_0 = \mathbb{Q}$ and $\mathfrak{Z}_1 = \{0\}$. Using any of the first two standard relations (2.3) or (4.4), one deduces $\mathfrak{Z}_p \cdot \mathfrak{Z}_{p'} \subset \mathfrak{Z}_{p+p'}$. This means that the \mathbb{Q} -vector subspace \mathfrak{Z} of \mathbb{R} spanned by all \mathfrak{Z}_p , $p \geq 0$, is a subalgebra of \mathbb{R} over \mathbb{Q} which is graded by the weight. From Conjecture 5.3 one deduces the following conjecture of Goncharov [G1]:

Conjecture 5.4. *As a \mathbb{Q} -algebra, \mathfrak{Z} is the direct sum of \mathfrak{Z}_p for $p \geq 0$.*

The dimension d_p of \mathfrak{Z}_p satisfies $d_0 = 1$, $d_1 = 0$, $d_2 = d_3 = 1$. The expected value for d_p is given by a conjecture of Zagier [Z]:

Conjecture 5.5. *For $p \geq 3$ we have*

$$d_p = d_{p-2} + d_{p-3}.$$

An interesting question is whether Conjecture 5.3 implies Conjecture 5.5. For this question as well as other related problems, see [É].

References

- [BR] Ball, K., Rivoal, T. – Irrationalité d’une infinité de valeurs de la fonction zêta aux entiers impairs. Submitted.
- [B³] Borwein, J.M., Bradley, D.M., Broadhurst, D.J. – Evaluations of k -fold Euler/Zagier sums: a compendium of results for arbitrary k . *Elec. J. Comb.* **4** (1997), N° 2, # R5, 21 pp.
- [B³L] Borwein, J.M., Bradley, D.M., Broadhurst, D.J., Linosek, P. – Special Values of Multiple Polylogarithms. *Trans. Amer. Math. Soc.*, (2000), to appear (35 pp.).
- [C] Cartier, P. – Sém. Bourbaki, Mars 2001.
- [É] Écalle, J. – Rapport sur les multizêtas et les sommes d’Euler. *Publ. Math. Orsay*, 11 pp., to appear.
- [Eu] Euler, L. – Meditationes circa singulare serierum genus. Leonhardi Euleri Opera Omnia, Series Prima XV, Commentationes Analyticae Vol. 2, 217–267; *Novi Comm. Acad. Sci. Petropol.*, **20** (1775), 140–186.
- [G1] Goncharov, A.B. – The double logarithms and Manin’s complex for modular curves. *Math. Research Letter* **4** (1997), n° 5, 6197–636.
- [G2] Goncharov, A.B. – Multiple polylogarithms, cyclotomy and modular complexes. *Math. Research Letter* **5** (1998), 497–516.
- [K] Kassel, C. – *Quantum Groups*. Graduate Texts in Math. **155**, Springer-Verlag, 1995.
- [MPH] Minh H. N., Petitot, M., Van Der Hoeven J. – Shuffle algebra and polylogarithms. Proc. of FPSAC’98, 10-th international Conference on *Formal Power Series and Algebraic Combinatorics*, June 98, Toronto.
- [R] Rivoal, T. – La fonction Zêta de Riemann prend une infinité de valeurs irrationnelles aux entiers impairs. *C. R. Acad. Sci. Paris* **331** (2000), 267–270.
<http://arXiv.org/abs/math.NT/0008051>
- [W] Waldschmidt, M. – Valeurs zêta multiples: une introduction. *J. Th. Nombres Bordeaux*, to appear (15 pp.).
<http://www.math.jussieu.fr/~miw/articles/ps/MZV.ps>
- [Z] Zagier, D. – Values of zeta functions and their applications. Proc. First European Congress of Mathematics, Vol. 2, Birkhauser, Boston (1994), 497–512.

<http://www.math.jussieu.fr/~miw/articles/ps/Chandigarh.ps>